

Permutation groups and transformation
semigroups:
3. Regularity and idempotent generation

Peter J. Cameron, University of St Andrews



Theoretical and Computational Algebra
July 2023

*You say I am repeating
Something I have said before. I shall say it again.
Shall I say it again?*

T. S. Eliot

Where are we going?

In the last lecture, we considered whether monoids of the form $\langle G, t \rangle$ are synchronizing for any non-permutation t . I didn't clearly explain why we just added one non-permutation; there are several reasons, for one the hope that we could proceed by induction.

Where are we going?

In the last lecture, we considered whether monoids of the form $\langle G, t \rangle$ are synchronizing for any non-permutation t . I didn't clearly explain why we just added one non-permutation; there are several reasons, for one the hope that we could proceed by induction.

It has to be said that synchronization is not the most significant property of finite semigroups. Semigroup theorists will propose other properties such as regularity and idempotent generation, which we will examine in this lecture. I will not justify their importance.

Where are we going?

In the last lecture, we considered whether monoids of the form $\langle G, t \rangle$ are synchronizing for any non-permutation t . I didn't clearly explain why we just added one non-permutation; there are several reasons, for one the hope that we could proceed by induction.

It has to be said that synchronization is not the most significant property of finite semigroups. Semigroup theorists will propose other properties such as regularity and idempotent generation, which we will examine in this lecture. I will not justify their importance.

A semigroup S is **regular** if every $a \in S$ has a **quasi-inverse** $b \in S$ such that $aba = a$ (and we may assume, without loss, that also $bab = b$). An **idempotent** is an element e satisfying $e^2 = e$; we say S is **idempotent-generated** if it is generated by its idempotents.

Where are we going?

In the last lecture, we considered whether monoids of the form $\langle G, t \rangle$ are synchronizing for any non-permutation t . I didn't clearly explain why we just added one non-permutation; there are several reasons, for one the hope that we could proceed by induction.

It has to be said that synchronization is not the most significant property of finite semigroups. Semigroup theorists will propose other properties such as regularity and idempotent generation, which we will examine in this lecture. I will not justify their importance.

A semigroup S is **regular** if every $a \in S$ has a **quasi-inverse** $b \in S$ such that $aba = a$ (and we may assume, without loss, that also $bab = b$). An **idempotent** is an element e satisfying $e^2 = e$; we say S is **idempotent-generated** if it is generated by its idempotents.

Note that if $aba = a$ then ab and ba are idempotents; so regularity implies the existence of idempotents, and idempotent-generation is stronger.

A nice theorem

The first thing to notice is that the analogue of our synchronization question was answered in 2011 by João Araújo, James Mitchell and Csaba Schneider:

A nice theorem

The first thing to notice is that the analogue of our synchronization question was answered in 2011 by João Araújo, James Mitchell and Csaba Schneider:

Theorem

Let G be a permutation group on Ω , with $|\Omega| = n$.

- ▶ *$\langle G, t \rangle$ is regular for any non-permutation t if and only if either G is $\text{Sym}(\Omega)$ or $\text{Alt}(\Omega)$, or G is one of nine specific permutation groups with $n = 5, 6, 7, 8$ or 9 .*

A nice theorem

The first thing to notice is that the analogue of our synchronization question was answered in 2011 by João Araújo, James Mitchell and Csaba Schneider:

Theorem

Let G be a permutation group on Ω , with $|\Omega| = n$.

- ▶ $\langle G, t \rangle$ is regular for any non-permutation t if and only if either G is $\text{Sym}(\Omega)$ or $\text{Alt}(\Omega)$, or G is one of nine specific permutation groups with $n = 5, 6, 7, 8$ or 9 .
- ▶ $\langle G, t \rangle \setminus G$ is idempotent-generated for any non-permutation t if and only if either G is $\text{Sym}(\Omega)$ or $\text{Alt}(\Omega)$, or G is one of three specific groups of degrees 5 or 6 .

A nice theorem

The first thing to notice is that the analogue of our synchronization question was answered in 2011 by João Araújo, James Mitchell and Csaba Schneider:

Theorem

Let G be a permutation group on Ω , with $|\Omega| = n$.

- ▶ $\langle G, t \rangle$ is regular for any non-permutation t if and only if either G is $\text{Sym}(\Omega)$ or $\text{Alt}(\Omega)$, or G is one of nine specific permutation groups with $n = 5, 6, 7, 8$ or 9 .
- ▶ $\langle G, t \rangle \setminus G$ is idempotent-generated for any non-permutation t if and only if either G is $\text{Sym}(\Omega)$ or $\text{Alt}(\Omega)$, or G is one of three specific groups of degrees 5 or 6 .

In the second statement, we remove elements of G since non-trivial permutations cannot be products of idempotents.

Set-transitive permutation groups

This result reminded me of a theorem with an interesting history. A permutation group G on Ω is said to be **set-transitive** if, for any two subsets of Ω of the same size, there is an element of G carrying the first to the second. In his book on permutation groups, Wielandt attributes the classification of these groups to Bercov. But also von Neumann and Morgenstern, in their pioneering book on game theory, stated the classification problem, and in the second edition asserted that it had been solved by Chevalley.

Set-transitive permutation groups

This result reminded me of a theorem with an interesting history. A permutation group G on Ω is said to be **set-transitive** if, for any two subsets of Ω of the same size, there is an element of G carrying the first to the second. In his book on permutation groups, Wielandt attributes the classification of these groups to Bercov. But also von Neumann and Morgenstern, in their pioneering book on game theory, stated the classification problem, and in the second edition asserted that it had been solved by Chevalley.

Theorem

Let G be a set-transitive permutation group on Ω , with $|\Omega| = n$. Then either G is $\text{Sym}(\Omega)$ or $\text{Alt}(\Omega)$, or G is one of four specific groups with $n = 5, 6$ or 9 .

We note that this theorem long predates CFSG.

A harder problem

So we need to make our original question more interesting, and the history of permutation groups gives a suggestion for how to do this.

A harder problem

So we need to make our original question more interesting, and the history of permutation groups gives a suggestion for how to do this.

In the first lecture, I discussed k -homogeneous permutation groups (those whose induced action on the set of k -element subsets of the domain is transitive). We saw that, for $2 \leq k \leq n/2$, these groups were shown to be k -transitive with known exceptions for $k = 2, 3$ or 4 , by the Livingstone–Wagner theorem and follow-up results of Kantor; then the CFSG allows us to determine the k -transitive groups.

A harder problem

So we need to make our original question more interesting, and the history of permutation groups gives a suggestion for how to do this.

In the first lecture, I discussed k -homogeneous permutation groups (those whose induced action on the set of k -element subsets of the domain is transitive). We saw that, for $2 \leq k \leq n/2$, these groups were shown to be k -transitive with known exceptions for $k = 2, 3$ or 4 , by the Livingstone–Wagner theorem and follow-up results of Kantor; then the CFSG allows us to determine the k -transitive groups.

In particular, the only k -transitive groups with $6 \leq k \leq |\Omega|/2$ are $\text{Sym}(\Omega)$ and $\text{Alt}(\Omega)$; for $k = 5$ there are additionally two **Mathieu groups** of degrees $n = 12$ and 24 , and for $k = 4$ two further Mathieu groups with $n = 11$ and 23 . For $k = 2$ or 3 there are infinitely many groups; but all are known.

In our situation . . .

So the question about “all subsets” was extended to the same question for “all subsets of size k ”. Similarly, in our problem, we can extend “all non-permutations” to “all transformations of rank k ”.

In our situation . . .

So the question about “all subsets” was extended to the same question for “all subsets of size k ”. Similarly, in our problem, we can extend “all non-permutations” to “all transformations of rank k ”.

So the first version reads:

Question

For which permutation groups G on Ω is it true that, for given k with $1 < k < n$,

- ▶ *if t is any map of rank k , then $\langle G, t \rangle$ is regular; or*

In our situation . . .

So the question about “all subsets” was extended to the same question for “all subsets of size k ”. Similarly, in our problem, we can extend “all non-permutations” to “all transformations of rank k ”.

So the first version reads:

Question

For which permutation groups G on Ω is it true that, for given k with $1 < k < n$,

- ▶ *if t is any map of rank k , then $\langle G, t \rangle$ is regular; or*
- ▶ *if t is any map of rank k , then $\langle G, t \rangle \setminus G$ is idempotent-generated?*

In our situation . . .

So the question about “all subsets” was extended to the same question for “all subsets of size k ”. Similarly, in our problem, we can extend “all non-permutations” to “all transformations of rank k ”.

So the first version reads:

Question

For which permutation groups G on Ω is it true that, for given k with $1 < k < n$,

- ▶ *if t is any map of rank k , then $\langle G, t \rangle$ is regular; or*
- ▶ *if t is any map of rank k , then $\langle G, t \rangle \setminus G$ is idempotent-generated?*

The basic strategy is to reformulate the hypotheses in a way which somehow resembles k -homogeneity.

The k -universal transversal property

A permutation group G on Ω is said to have the **k -universal transversal property**, or **k -ut** for short (where $1 \leq k \leq n - 1$) if, given any k -subset A and k -partition P of Ω , there is an element $g \in G$ mapping A to a transversal for P .

Proposition

The permutation group G has the property that $\langle G, t \rangle$ is regular for all maps t of rank k if and only if G has the k -universal transversal property.

This is exactly the kind of reformulation we are looking for, and its study involves interesting combinatorics and also reveals parallels to the Livingstone–Wagner theorem.

The k -universal transversal property

A permutation group G on Ω is said to have the **k -universal transversal property**, or **k -ut** for short (where $1 \leq k \leq n - 1$) if, given any k -subset A and k -partition P of Ω , there is an element $g \in G$ mapping A to a transversal for P .

Proposition

The permutation group G has the property that $\langle G, t \rangle$ is regular for all maps t of rank k if and only if G has the k -universal transversal property.

This is exactly the kind of reformulation we are looking for, and its study involves interesting combinatorics and also reveals parallels to the Livingstone–Wagner theorem.

But we have to take quite a journey to get there.

k -ut and regularity

It is not hard to prove that, if G has the k -ut and t has rank k , then t has a quasi-inverse in $S = \langle G, t \rangle$.

k -ut and regularity

It is not hard to prove that, if G has the k -ut and t has rank k , then t has a quasi-inverse in $S = \langle G, t \rangle$.

Take $g \in G$ mapping $\text{Im}(t)$ to a transversal for $\text{Ker}(t)$. Then gt induces a permutation of $\text{Im}(t)$, and so some power of it (say $(gt)^m$) is the identity on $\text{Im}(t)$; then $t(gt)^m = t$. (Note in passing that $(gt)^m$ is an idempotent.)

k -ut and regularity

It is not hard to prove that, if G has the k -ut and t has rank k , then t has a quasi-inverse in $S = \langle G, t \rangle$.

Take $g \in G$ mapping $\text{Im}(t)$ to a transversal for $\text{Ker}(t)$. Then gt induces a permutation of $\text{Im}(t)$, and so some power of it (say $(gt)^m$) is the identity on $\text{Im}(t)$; then $t(gt)^m = t$. (Note in passing that $(gt)^m$ is an idempotent.)

So if we knew that every element of rank at most k in $\langle G, t \rangle$ had a quasi-inverse, we would be done. In other words, we have to prove that k -ut implies $(k-1)$ -ut (for $k \leq n/2$), and use induction.

Going down

Livingstone and Wagner are our guides. Their first theorem stated that, if $k \leq n/2$ and G is k -homogeneous, then G is $(k - 1)$ -homogeneous.

Going down

Livingstone and Wagner are our guides. Their first theorem stated that, if $k \leq n/2$ and G is k -homogeneous, then G is $(k - 1)$ -homogeneous.

Their proof used facts about character theory of symmetric groups. Soon after, Wielandt found a number-theoretic proof and Kantor a combinatorial proof.

Going down

Livingstone and Wagner are our guides. Their first theorem stated that, if $k \leq n/2$ and G is k -homogeneous, then G is $(k - 1)$ -homogeneous.

Their proof used facts about character theory of symmetric groups. Soon after, Wielandt found a number-theoretic proof and Kantor a combinatorial proof.

We need to use consequences of CFSG.

Going down

Livingstone and Wagner are our guides. Their first theorem stated that, if $k \leq n/2$ and G is k -homogeneous, then G is $(k - 1)$ -homogeneous.

Their proof used facts about character theory of symmetric groups. Soon after, Wielandt found a number-theoretic proof and Kantor a combinatorial proof.

We need to use consequences of CFSG.

I make one observation here.

Proposition

The 2-ut property is equivalent to primitivity.

Going down

Livingstone and Wagner are our guides. Their first theorem stated that, if $k \leq n/2$ and G is k -homogeneous, then G is $(k - 1)$ -homogeneous.

Their proof used facts about character theory of symmetric groups. Soon after, Wielandt found a number-theoretic proof and Kantor a combinatorial proof.

We need to use consequences of CFSG.

I make one observation here.

Proposition

The 2-ut property is equivalent to primitivity.

For let Γ be a G -invariant graph, and A an edge. If G has 2-ut, then for any 2-part partition, there is an image of A under G having one point in each part. Hence Γ is connected, and the result follows from Higman's theorem.

Going down, 2

We say that a permutation group G is (l, k) homogeneous (for $l < k$) if given any l -set A and k -set B , G contains an element mapping A to a subset of B .

Going down, 2

We say that a permutation group G is (l, k) homogeneous (for $l < k$) if given any l -set A and k -set B , G contains an element mapping A to a subset of B .

Now k -ut implies $(k - 1, k)$ -homogeneous. For let B be a k -set and P a partition whose parts are the singletons of a $(k - 1)$ -set A and a single part containing all the rest; let g map A to a transversal to P . Then g^{-1} maps B to a subset of A .

Going down, 2

We say that a permutation group G is (l, k) homogeneous (for $l < k$) if given any l -set A and k -set B , G contains an element mapping A to a subset of B .

Now k -ut implies $(k - 1, k)$ -homogeneous. For let B be a k -set and P a partition whose parts are the singletons of a $(k - 1)$ -set A and a single part containing all the rest; let g map A to a transversal to P . Then g^{-1} maps B to a subset of A .

Now we classify $(k - 1, k)$ -homogeneous groups; use combinatorics (including Ramsey's theorem) to show that such a group is 2-homogeneous (at least for $k \geq 3$) with a few known exceptions, and then appeal to the classification of such groups. Once they are all known, it is only necessary to check that they all have $(k - 1)$ -ut, and we are done.

Comments

This theorem was proved by João and me and published in 2016. The proof involved an elaborate mixture of combinatorics, group theory, and computation.

Comments

This theorem was proved by João and me and published in 2016. The proof involved an elaborate mixture of combinatorics, group theory, and computation.

We also gave a partial classification of k -ut groups for $2 < k \leq n/2$. This was subsequently pushed a little further with the help of Wolfram Bentz, but the job is not quite finished yet.

Comments

This theorem was proved by João and me and published in 2016. The proof involved an elaborate mixture of combinatorics, group theory, and computation.

We also gave a partial classification of k -ut groups for $2 < k \leq n/2$. This was subsequently pushed a little further with the help of Wolfram Bentz, but the job is not quite finished yet.

However, if we regard that answer as satisfactory, we could ask, what next? We can refine the question further by asking about groups G for which $\langle G, t \rangle$ is regular for just some of the maps of rank k . One such weakening is the k -et property, to which I turn next.

k -existential transversal property

A permutation group G has the k -existential transversal property, or k -et for short, if there is a k -subset A of the domain (called a witness) such that, for any map t with $\text{Im}(t) = A$, the semigroup $\langle G, t \rangle$ is regular.

k -existential transversal property

A permutation group G has the k -existential transversal property, or k -et for short, if there is a k -subset A of the domain (called a witness) such that, for any map t with $\text{Im}(t) = A$, the semigroup $\langle G, t \rangle$ is regular.

This is considerably weaker than k -ut, but we have a similarly almost satisfactory result, including a nearly complete classification for $k \geq 4$.

k -existential transversal property

A permutation group G has the **k -existential transversal property**, or k -et for short, if there is a k -subset A of the domain (called a witness) such that, for any map t with $\text{Im}(t) = A$, the semigroup $\langle G, t \rangle$ is regular.

This is considerably weaker than k -ut, but we have a similarly almost satisfactory result, including a nearly complete classification for $k \geq 4$.

One curious feature is that the going-down result almost holds:

Theorem

Suppose that G is a permutation group of degree n , which has the k -et property for $2 \leq k \leq n/2$. Then either G has the $(k-1)$ -et property, or G is one of two affine groups of degree 16, with $k = 6$.

k -existential transversal property

A permutation group G has the **k -existential transversal property**, or k -et for short, if there is a k -subset A of the domain (called a witness) such that, for any map t with $\text{Im}(t) = A$, the semigroup $\langle G, t \rangle$ is regular.

This is considerably weaker than k -ut, but we have a similarly almost satisfactory result, including a nearly complete classification for $k \geq 4$.

One curious feature is that the going-down result almost holds:

Theorem

Suppose that G is a permutation group of degree n , which has the k -et property for $2 \leq k \leq n/2$. Then either G has the $(k-1)$ -et property, or G is one of two affine groups of degree 16, with $k = 6$.

There are other strange beasts too: for example, the Higman–Sims group and its automorphism group (with $n = 100$) are the only groups with 4-et and $n \geq 8$ which are not 2-homogeneous.

2-et

We saw that the 2-ut property is equivalent to primitivity. So the 2-et property is a weakening of primitivity.

2-et

We saw that the 2-ut property is equivalent to primitivity. So the 2-et property is a weakening of primitivity.

Higman's theorem says that a group is primitive if and only if all its orbital graphs are connected. In the other direction, we say that G is **fully imprimitive** if all its orbital graphs are disconnected.

2-et

We saw that the 2-ut property is equivalent to primitivity. So the 2-et property is a weakening of primitivity.

Higman's theorem says that a group is primitive if and only if all its orbital graphs are connected. In the other direction, we say that G is **fully imprimitive** if all its orbital graphs are disconnected.

Proposition

A permutation group G has the 2-et property if and only if it is not fully imprimitive.

2-et

We saw that the 2-ut property is equivalent to primitivity. So the 2-et property is a weakening of primitivity.

Higman's theorem says that a group is primitive if and only if all its orbital graphs are connected. In the other direction, we say that G is **fully imprimitive** if all its orbital graphs are disconnected.

Proposition

A permutation group G has the 2-et property if and only if it is not fully imprimitive.

In particular, a regular permutation group G (that is, acting on itself by right multiplication) is fully imprimitive if and only if it is not cyclic.

Variants

I have been discussing submonoids of the full transformation monoid T_n generated by a permutation group and one non-permutation.

Variants

I have been discussing submonoids of the full transformation monoid T_n generated by a permutation group and one non-permutation.

But we can play this game in other arenas. For example, the **symmetric inverse semigroup** I_n of all **partial permutations** (bijective maps between subsets of Ω). More generally, $PT(n)$ denotes the semigroup of **partial transformations** of Ω .

Variants

I have been discussing submonoids of the full transformation monoid T_n generated by a permutation group and one non-permutation.

But we can play this game in other arenas. For example, the **symmetric inverse semigroup** I_n of all **partial permutations** (bijective maps between subsets of Ω). More generally, $PT(n)$ denotes the semigroup of **partial transformations** of Ω .

One could also play the game in other famous semigroups such as the **Brauer monoid** and the **partition monoid**, or linear versions in the **general linear group and semigroup**. This has yet to be done.

(k, l) -ut

Consider the partial transformation monoid. We want to know, given k, l with $k < l$, which permutation groups G have the property that, for any map t with domain of size l and image of size k , the semigroup $\langle G, t \rangle$ is regular.

(k, l) -ut

Consider the partial transformation monoid. We want to know, given k, l with $k < l$, which permutation groups G have the property that, for any map t with domain of size l and image of size k , the semigroup $\langle G, t \rangle$ is regular.

There is a combinatorial property of G , the (k, l) -ut property, which guarantees this (if $k \leq n/2$): G has the (k, l) property if, given a k -set A , an l -set B , and a k -partition of B , there is an element $g \in G$ mapping A to a transversal for B .

(k, l) -ut

Consider the partial transformation monoid. We want to know, given k, l with $k < l$, which permutation groups G have the property that, for any map t with domain of size l and image of size k , the semigroup $\langle G, t \rangle$ is regular.

There is a combinatorial property of G , the (k, l) -ut property, which guarantees this (if $k \leq n/2$): G has the (k, l) property if, given a k -set A , an l -set B , and a k -partition of B , there is an element $g \in G$ mapping A to a transversal for B .

Theorem

- ▶ (k, n) -ut is equivalent to k -ut, and (k, k) -ut to k -homogeneity.

(k, l) -ut

Consider the partial transformation monoid. We want to know, given k, l with $k < l$, which permutation groups G have the property that, for any map t with domain of size l and image of size k , the semigroup $\langle G, t \rangle$ is regular.

There is a combinatorial property of G , the (k, l) -ut property, which guarantees this (if $k \leq n/2$): G has the (k, l) property if, given a k -set A , an l -set B , and a k -partition of B , there is an element $g \in G$ mapping A to a transversal for B .

Theorem

- ▶ (k, n) -ut is equivalent to k -ut, and (k, k) -ut to k -homogeneity.
- ▶ The (k, l) -ut property becomes weaker as l increases from k to n ; so, if G has k -ut, then there is a threshold $t(G, k)$ such that G has (k, l) -ut if and only if $t(G, k) \leq l \leq n$.

(k, l) -ut

Consider the partial transformation monoid. We want to know, given k, l with $k < l$, which permutation groups G have the property that, for any map t with domain of size l and image of size k , the semigroup $\langle G, t \rangle$ is regular.

There is a combinatorial property of G , the (k, l) -ut property, which guarantees this (if $k \leq n/2$): G has the (k, l) property if, given a k -set A , an l -set B , and a k -partition of B , there is an element $g \in G$ mapping A to a transversal for B .

Theorem

- ▶ (k, n) -ut is equivalent to k -ut, and (k, k) -ut to k -homogeneity.
- ▶ The (k, l) -ut property becomes weaker as l increases from k to n ; so, if G has k -ut, then there is a threshold $t(G, k)$ such that G has (k, l) -ut if and only if $t(G, k) \leq l \leq n$.

The value of $T(G, k)$ is known for most groups G with k -ut.

Idempotent generation

As we saw, the k -ut property guarantees that $\langle G, t \rangle$ contains idempotents when $\text{rank}(t) = k$. Asking for it to be generated by idempotents is a stronger condition on G which we call **k -idempotent generated** (for short, k -id).

Idempotent generation

As we saw, the k -ut property guarantees that $\langle G, t \rangle$ contains idempotents when $\text{rank}(t) = k$. Asking for it to be generated by idempotents is a stronger condition on G which we call **k -idempotent generated** (for short, k -id).

Since we have a reasonably good, if not complete, account of k -ut groups, we should be able to read off the k -id groups, yes?

Idempotent generation

As we saw, the k -ut property guarantees that $\langle G, t \rangle$ contains idempotents when $\text{rank}(t) = k$. Asking for it to be generated by idempotents is a stronger condition on G which we call **k -idempotent generated** (for short, k -id).

Since we have a reasonably good, if not complete, account of k -ut groups, we should be able to read off the k -id groups, yes? In fact the problem is that, apart from $k = 2$, we don't understand these groups. We have a condition called **strong k -ut**, and k -id is sandwiched between k -ut and strong k -ut, but we don't have a purely combinatorial or group-theoretic condition for k -id.

The Road Closure Property

Let G be a transitive permutation group on Ω .

The Road Closure Property

Let G be a transitive permutation group on Ω .

Recall: an **orbital graph** for G is a graph obtained by taking as edge set an orbit of G on 2-element subsets; by **Higman's Theorem**, G is primitive if and only if every orbital graph for G is connected.

The Road Closure Property

Let G be a transitive permutation group on Ω .

Recall: an **orbital graph** for G is a graph obtained by taking as edge set an orbit of G on 2-element subsets; by **Higman's Theorem**, G is primitive if and only if every orbital graph for G is connected.

The **Road Closure Property** is a strengthening of this. G has the Road Closure Property (RCP for short) if, whenever E is an orbit of G on 2-sets, and B a proper block of imprimitivity for the action of G on E , the graph with edge set $E \setminus B$ is connected. In other words, if all the roads in a block are closed for roadworks, it is still possible to get around the network.

The Road Closure Property

Let G be a transitive permutation group on Ω .

Recall: an **orbital graph** for G is a graph obtained by taking as edge set an orbit of G on 2-element subsets; by **Higman's Theorem**, G is primitive if and only if every orbital graph for G is connected.

The **Road Closure Property** is a strengthening of this. G has the Road Closure Property (RCP for short) if, whenever E is an orbit of G on 2-sets, and B a proper block of imprimitivity for the action of G on E , the graph with edge set $E \setminus B$ is connected. In other words, if all the roads in a block are closed for roadworks, it is still possible to get around the network.

Theorem

The permutation group G has the 2-id property if and only if it has the Road Closure Property.

Examples

Clearly a group with the Road Closure Property is primitive. Moreover, it is basic. For a Hamming graph $H(m, q)$ on Ω gives it the structure of an m -dimensional cube (with q points on each edge); the m directions in the cube form a system of imprimitivity. If we delete all edges in one direction, the graph falls apart into m layers.

Examples

Clearly a group with the Road Closure Property is primitive. Moreover, it is basic. For a Hamming graph $H(m, q)$ on Ω gives it the structure of an m -dimensional cube (with q points on each edge); the m directions in the cube form a system of imprimitivity. If we delete all edges in one direction, the graph falls apart into m layers.

Another family is exemplified by the group of collineations and dualities of a projective plane over a finite field, acting on the set of flags (incident point-line pairs) in the plane; edges consist of pairs of flags sharing a point or a line. There is a natural system of two blocks of edges; if we delete, say, all edges consisting of pairs of flags sharing a point, then we cannot move from flags through one point to flags through another. Many other incidence structures provide similar examples.

Examples

Clearly a group with the Road Closure Property is primitive. Moreover, it is basic. For a Hamming graph $H(m, q)$ on Ω gives it the structure of an m -dimensional cube (with q points on each edge); the m directions in the cube form a system of imprimitivity. If we delete all edges in one direction, the graph falls apart into m layers.

Another family is exemplified by the group of collineations and dualities of a projective plane over a finite field, acting on the set of flags (incident point-line pairs) in the plane; edges consist of pairs of flags sharing a point or a line. There is a natural system of two blocks of edges; if we delete, say, all edges consisting of pairs of flags sharing a point, then we cannot move from flags through one point to flags through another. Many other incidence structures provide similar examples. There are a few other examples too: Pablo Spiga found what is now known to be the smallest example with more than two blocks, the group $\text{PTU}(3, 5)$ acting on 6000 points.

Comments

We conjecture that the examples we have seen are typical: that is, a basic primitive group which fails the RCP has two or three blocks of imprimitivity on the edges of some orbital graph. However, we are still some way from a proof.

Comments

We conjecture that the examples we have seen are typical: that is, a basic primitive group which fails the RCP has two or three blocks of imprimitivity on the edges of some orbital graph.

However, we are still some way from a proof.

The RCP, like synchronization, is a strengthening of primitivity. It is tempting to think that there might be a connection between these two properties, but none is currently known or suspected.

References

- ▶ J. Araújo, J. P. Araújo, W. Bentz, P. J. Cameron and P. Spiga, A transversal property for permutation groups motivated by partial transformations, *J. Algebra* **573** (2021), 741–759.
- ▶ João Araújo, Wolfram Bentz and Peter J. Cameron, The existential transversal property: a generalization of homogeneity and its impact on semigroups, *Trans. Amer. Math. Soc.* **374** (2021), 1155–1195.
- ▶ João Araújo and Peter J. Cameron, Two generalizations of homogeneity in groups with applications to regular semigroups, *Trans. Amer. Math. Soc.* **368** (2016), 1159–1188.
- ▶ João Araújo and Peter J. Cameron, Primitive groups, road closures and idempotent generation, arXiv 1611.08233.