

Semper Abstracta?

Peter J. Cameron
University of St Andrews
(Emeritus)



Theoretical and Computational Algebra
Évora, July 2025

João Araújo



João Araújo



I am grateful to João for many things, including

João Araújo



I am grateful to João for many things, including

- ▶ introducing me to many beautiful parts of Portugal;

João Araújo



I am grateful to João for many things, including

- ▶ introducing me to many beautiful parts of Portugal;
- ▶ introducing me to many beautiful mathematical problems;

João Araújo



I am grateful to João for many things, including

- ▶ introducing me to many beautiful parts of Portugal;
- ▶ introducing me to many beautiful mathematical problems;
- ▶ masterminding this beautiful conference.

Semper Abstracta?

The motto on my title slide announces a good maxim for mathematicians. Our subject derives much of its power from its abstract nature; the results we prove can be applied to many different areas.

Semper Abstracta?

The motto on my title slide announces a good maxim for mathematicians. Our subject derives much of its power from its abstract nature; the results we prove can be applied to many different areas.

But it is not quite so simple.

Semper Abstracta?

The putto on my title slide announces a good maxim for mathematicians. Our subject derives much of its power from its abstract nature; the results we prove can be applied to many different areas.

But it is not quite so simple.

There is a line between the abstract and the concrete, and any particular piece of mathematics can be positioned somewhere on that line.

Semper Abstracta?

The putto on my title slide announces a good maxim for mathematicians. Our subject derives much of its power from its abstract nature; the results we prove can be applied to many different areas.

But it is not quite so simple.

There is a line between the abstract and the concrete, and any particular piece of mathematics can be positioned somewhere on that line.

And more seriously, there may be room for considerable disagreement about where to put it.

Semper Abstracta?

The putto on my title slide announces a good maxim for mathematicians. Our subject derives much of its power from its abstract nature; the results we prove can be applied to many different areas.

But it is not quite so simple.

There is a line between the abstract and the concrete, and any particular piece of mathematics can be positioned somewhere on that line.

And more seriously, there may be room for considerable disagreement about where to put it.

In the interview for my postdoctoral fellowship in Oxford in 1971, I was asked about some discussions I had had with Don Taylor about a configuration of points in 23-dimensional Euclidean space (a subset of the Leech lattice). We thought that this was extremely concrete. But the committee (mostly non-mathematicians) were so impressed that somebody could think in 23 dimensions that I got the position.

Concrete or abstract?

A mathematician, like a painter or poet, is a maker of patterns ... Beauty is the first test; there is no place in the world for ugly mathematics.

G. H. Hardy, *A Mathematician's Apology*

Concrete or abstract?

A mathematician, like a painter or poet, is a maker of patterns ... Beauty is the first test; there is no place in the world for ugly mathematics.

G. H. Hardy, *A Mathematician's Apology*

... the heart of the poem is given to the concrete, and it is in this direction that the poem goes. This is the diametric opposite of the ping-pong of mathematics, in which the last shot is always towards the abstract.

Ron Aharoni, *Mathematics, Poetry and Beauty*

Concrete or abstract?

A mathematician, like a painter or poet, is a maker of patterns ... Beauty is the first test; there is no place in the world for ugly mathematics.

G. H. Hardy, *A Mathematician's Apology*

... the heart of the poem is given to the concrete, and it is in this direction that the poem goes. This is the diametric opposite of the ping-pong of mathematics, in which the last shot is always towards the abstract.

Ron Aharoni, *Mathematics, Poetry and Beauty*

I am not so sure, however.

How to teach algebra

Groups, then rings; or rings, then groups?

How to teach algebra

Groups, then rings; or rings, then groups?

I have always taken the view that fewer axioms don't mean easier to understand for a beginner.

How to teach algebra

Groups, then rings; or rings, then groups?

I have always taken the view that fewer axioms don't mean easier to understand for a beginner.

Rings are fairly concrete objects, since there is a prototype or exemplar, the ring of integers, with which all our students are familiar. Once they have mastered an axiomatic system with the integers to guide them, we can make things more abstract, by stripping away axioms, to study groups, and then maybe semigroups.

How to teach algebra

Groups, then rings; or rings, then groups?

I have always taken the view that fewer axioms don't mean easier to understand for a beginner.

Rings are fairly concrete objects, since there is a prototype or exemplar, the ring of integers, with which all our students are familiar. Once they have mastered an axiomatic system with the integers to guide them, we can make things more abstract, by stripping away axioms, to study groups, and then maybe semigroups.

If you begin with groups, there is no natural exemplar.

Abstract or concrete? Road Closure

Abstract

A permutation group G on Ω has the **Road Closure Property** if no orbital graph for G can be disconnected by deleting the edges in a block of imprimitivity for G in its action on edges.

Concrete



Concrete or abstract?

This question is related to the most important question in the philosophy of mathematics: “Discovered or invented?”. These questions are not equivalent, but a definitive answer to one would have implications for the other.

Concrete or abstract?

This question is related to the most important question in the philosophy of mathematics: “Discovered or invented?”. These questions are not equivalent, but a definitive answer to one would have implications for the other.

Mathematics becomes more concrete as it moves from mathematician to physicist, engineer, economist, etc. Also as it moves from general to particular. For example, “A group having exactly 59 subgroups” is surely more abstract than “The alternating group on 5 letters”, although they are exactly equivalent.

God said, "Let there be light," and light came into existence, perhaps in a rather undefined form; it takes its specifics (frequencies and intensities) from the objects emitting or reflecting it. Then it can create the visible universe in the eye of the beholder.

God said, "Let there be light," and light came into existence, perhaps in a rather undefined form; it takes its specifics (frequencies and intensities) from the objects emitting or reflecting it. Then it can create the visible universe in the eye of the beholder.

A mathematician, at the start of a lecture, says "Let G be a group," and a group is called into existence in the minds of the listeners, in a rather undefined form at first; but as the lecture proceeds, the group takes on more specific properties, and perhaps by the end of the lecture it has become a single group, or a narrowly specified class of groups.

Talk outline

My talk will consist of a series of examples to illustrate the points. Questions are welcome, so it may be that I don't cover everything. The topics will be

Talk outline

My talk will consist of a series of examples to illustrate the points. Questions are welcome, so it may be that I don't cover everything. The topics will be

- ▶ the random graph and Zermelo–Fraenkel set theory,

Talk outline

My talk will consist of a series of examples to illustrate the points. Questions are welcome, so it may be that I don't cover everything. The topics will be

- ▶ the random graph and Zermelo–Fraenkel set theory,
- ▶ a problem about derangements,

Talk outline

My talk will consist of a series of examples to illustrate the points. Questions are welcome, so it may be that I don't cover everything. The topics will be

- ▶ the random graph and Zermelo–Fraenkel set theory,
- ▶ a problem about derangements,
- ▶ the ADE Coxeter–Dynkin diagrams.

Talk outline

My talk will consist of a series of examples to illustrate the points. Questions are welcome, so it may be that I don't cover everything. The topics will be

- ▶ the random graph and Zermelo–Fraenkel set theory,
- ▶ a problem about derangements,
- ▶ the ADE Coxeter–Dynkin diagrams.

I will start each new section by showing you a picture of Évora, so if you have lost the thread you can come back in at that point.



The random graph and ZF set theory

Zermelo–Fraenkel set theory is highly abstract, since it is an axiom system for set theory which is the most usual basis for all of mathematics.

The random graph and ZF set theory

Zermelo–Fraenkel set theory is highly abstract, since it is an axiom system for set theory which is the most usual basis for all of mathematics.

The countable random graph was a discovery of Erdős and Rényi. There is a graph R with the following remarkable property:

If you take a countable set of vertices, and for each pair of vertices, choose independently at random (for example, by tossing a coin) whether this pair are joined by an edge or not. Let X be the resulting random graph. Then X is almost surely isomorphic to R (that is, $\mathbb{P}(X \cong R) = 1$).

The random graph and ZF set theory

Zermelo–Fraenkel set theory is highly abstract, since it is an axiom system for set theory which is the most usual basis for all of mathematics.

The countable random graph was a discovery of Erdős and Rényi. There is a graph R with the following remarkable property:

If you take a countable set of vertices, and for each pair of vertices, choose independently at random (for example, by tossing a coin) whether this pair are joined by an edge or not. Let X be the resulting random graph. Then X is almost surely isomorphic to R (that is, $\mathbb{P}(X \cong R) = 1$).

This already much more concrete than ZF set theory (although it is a non-constructive existence proof of R , a fairly abstract idea). But it can be made more concrete still. Independently of Erdős and Rényi, Rado constructed a graph R as on the next slide.

Rado's construction

Rado defined his graph as follows. Let \mathbb{N} be the set of natural numbers (including 0). Rado's graph has vertex set \mathbb{N} . The rule for joining i and j is as follows. Suppose that $i < j$ (as we can always do). Now express j in base 2. If the i th digit is 1, we join i to j , otherwise not.

Rado's construction

Rado defined his graph as follows. Let \mathbb{N} be the set of natural numbers (including 0). Rado's graph has vertex set \mathbb{N} . The rule for joining i and j is as follows. Suppose that $i < j$ (as we can always do). Now express j in base 2. If the i th digit is 1, we join i to j , otherwise not.

Thus, for example, 0 is joined to all odd numbers; 1 is joined to 0 and to all numbers congruent to 2 or 3 (mod 4).

Rado's construction

Rado defined his graph as follows. Let \mathbb{N} be the set of natural numbers (including 0). Rado's graph has vertex set \mathbb{N} . The rule for joining i and j is as follows. Suppose that $i < j$ (as we can always do). Now express j in base 2. If the i th digit is 1, we join i to j , otherwise not.

Thus, for example, 0 is joined to all odd numbers; 1 is joined to 0 and to all numbers congruent to 2 or 3 (mod 4).

The connection is as follows:

Rado's construction

Rado defined his graph as follows. Let \mathbb{N} be the set of natural numbers (including 0). Rado's graph has vertex set \mathbb{N} . The rule for joining i and j is as follows. Suppose that $i < j$ (as we can always do). Now express j in base 2. If the i th digit is 1, we join i to j , otherwise not.

Thus, for example, 0 is joined to all odd numbers; 1 is joined to 0 and to all numbers congruent to 2 or 3 (mod 4).

The connection is as follows:

Theorem

Rado's graph is isomorphic to R .

The connection

According to the **downward Löwenheim–Skolem Theorem**, if a set of first-order sentences in a countable language has a model, then it has a countable model. So, if ZF is consistent (as we all hope!), it has a countable model. This is despite the fact that a theorem of ZF is the existence of uncountable sets! This is the **Skolem paradox**, which I am not going to resolve here.

The connection

According to the **downward Löwenheim–Skolem Theorem**, if a set of first-order sentences in a countable language has a model, then it has a countable model. So, if ZF is consistent (as we all hope!), it has a countable model. This is despite the fact that a theorem of ZF is the existence of uncountable sets! This is the **Skolem paradox**, which I am not going to resolve here.

Now the language of ZF has a single binary relation, \in (membership). So a model is a directed graph satisfying a rather complicated set of axioms.

The connection

According to the **downward Löwenheim–Skolem Theorem**, if a set of first-order sentences in a countable language has a model, then it has a countable model. So, if ZF is consistent (as we all hope!), it has a countable model. This is despite the fact that a theorem of ZF is the existence of uncountable sets! This is the **Skolem paradox**, which I am not going to resolve here.

Now the language of ZF has a single binary relation, \in (membership). So a model is a directed graph satisfying a rather complicated set of axioms.

Theorem

If we take a countable model of ZF, and ignore the directions on the edges, we obtain the random graph R .

The connection

According to the **downward Löwenheim–Skolem Theorem**, if a set of first-order sentences in a countable language has a model, then it has a countable model. So, if ZF is consistent (as we all hope!), it has a countable model. This is despite the fact that a theorem of ZF is the existence of uncountable sets! This is the **Skolem paradox**, which I am not going to resolve here.

Now the language of ZF has a single binary relation, \in (membership). So a model is a directed graph satisfying a rather complicated set of axioms.

Theorem

If we take a countable model of ZF, and ignore the directions on the edges, we obtain the random graph R .

So the abstract touches the very concrete (Rado's construction).

There is more!

The proof of the last theorem uses by no means all of the ZF axioms. The most crucial axiom is the **Axiom of Foundation**. Other axioms, such as the axioms of Infinity and Choice are not required.

There is more!

The proof of the last theorem uses by no means all of the ZF axioms. The most crucial axiom is the **Axiom of Foundation**. Other axioms, such as the axioms of Infinity and Choice are not required.

In particular, if we take Rado's graph, and direct the edges from smaller to larger, we obtain a model of **hereditarily finite set theory** (obtained by replacing the Axiom of Infinity by its negation): all sets are finite, and all their members are finite, and so on all the way down.

There is more!

The proof of the last theorem uses by no means all of the ZF axioms. The most crucial axiom is the **Axiom of Foundation**. Other axioms, such as the axioms of Infinity and Choice are not required.

In particular, if we take Rado's graph, and direct the edges from smaller to larger, we obtain a model of **hereditarily finite set theory** (obtained by replacing the Axiom of Infinity by its negation): all sets are finite, and all their members are finite, and so on all the way down.

The sequel to this story is that I had a project student at St Andrews, Bea Adam-Day, who was fascinated by this story, and decided to do a project on what graphs are obtained if we replace the crucial Axiom of Foundation (forbidding infinite descending chains under membership) by the so-called **Anti-Foundation Axiom**. She did a good project, and then during her PhD at the University of Leeds, she and two fellow students essentially solved the problem completely.



Inverse group theory

This is an ongoing project with João Araújo, Francesco Matucci, and others.

Let F be an operation taking groups to groups (*not* necessarily a functor). The problem of inverse group theory is:

Given a group H , does there exist a group G such that $F(G) = H$? If so, can we describe all such groups?

Inverse group theory

This is an ongoing project with João Araújo, Francesco Matucci, and others.

Let F be an operation taking groups to groups (*not* necessarily a functor). The problem of inverse group theory is:

Given a group H , does there exist a group G such that $F(G) = H$? If so, can we describe all such groups?

Example Suppose that $F(G)$ is the derived group G' (the subgroup generated by commutators). A group G with $G' = H$ is called an **integral** of H .

Inverse group theory

This is an ongoing project with João Araújo, Francesco Matucci, and others.

Let F be an operation taking groups to groups (*not* necessarily a functor). The problem of inverse group theory is:

Given a group H , does there exist a group G such that $F(G) = H$? If so, can we describe all such groups?

Example Suppose that $F(G)$ is the derived group G' (the subgroup generated by commutators). A group G with $G' = H$ is called an **integral** of H .

Scandalously, we do not even know whether the computational problem of deciding whether a finite group is integrable is even decidable, let alone have a good estimate for its complexity!

Frobenius groups

I have always felt that permutation groups are more “concrete” than abstract groups. The purpose of a group is to act on something, and I believe that there are things to be learned from the interaction of the group with its operand.

Frobenius groups

I have always felt that permutation groups are more “concrete” than abstract groups. The purpose of a group is to act on something, and I believe that there are things to be learned from the interaction of the group with its operand.

One of the topics in inverse group theory arises from **Frobenius’ theorem**, which can be stated in two ways, which are entirely equivalent, but one is more concrete than the other.

Frobenius groups

I have always felt that permutation groups are more “concrete” than abstract groups. The purpose of a group is to act on something, and I believe that there are things to be learned from the interaction of the group with its operand.

One of the topics in inverse group theory arises from **Frobenius’ theorem**, which can be stated in two ways, which are entirely equivalent, but one is more concrete than the other.

Theorem

- ▶ *Let G be a finite transitive permutation group. Suppose that the stabiliser of any point is non-trivial, but the stabiliser of any two points is trivial. Then the identity and the derangements (fixed-point-free elements) in G form a regular normal subgroup, and the point stabilisers are complements.*

Frobenius groups

I have always felt that permutation groups are more “concrete” than abstract groups. The purpose of a group is to act on something, and I believe that there are things to be learned from the interaction of the group with its operand.

One of the topics in inverse group theory arises from **Frobenius’ theorem**, which can be stated in two ways, which are entirely equivalent, but one is more concrete than the other.

Theorem

- ▶ *Let G be a finite transitive permutation group. Suppose that the stabiliser of any point is non-trivial, but the stabiliser of any two points is trivial. Then the identity and the derangements (fixed-point-free elements) in G form a regular normal subgroup, and the point stabilisers are complements.*
- ▶ *Let G be a finite group with a subgroup H such that $H \neq 1$ but $H \cap H^g = 1$ for all $g \notin H$. Then H has a normal complement.*

The derangement subgroup

A few years ago, inspired by work of H. Zantema in number theory, Rosemary Bailey, Michael Giudici, Gordon Royle and I looked at the more general question. Let G be a finite transitive permutation group, and let $D(G)$ be the subgroup generated by the derangements in G . Which groups can arise as $G/D(G)$? (This fits well our paradigm of inverse group theory.)

The derangement subgroup

A few years ago, inspired by work of H. Zantema in number theory, Rosemary Bailey, Michael Giudici, Gordon Royle and I looked at the more general question. Let G be a finite transitive permutation group, and let $D(G)$ be the subgroup generated by the derangements in G . Which groups can arise as $G/D(G)$?

(This fits well our paradigm of inverse group theory.)

We found that the vast majority of transitive groups have $D(G) = G$. Of course, if G is a Frobenius group, then $D(G)$ is the Frobenius kernel, and the quotient is the Frobenius complement. The structure of Frobenius complements is known very precisely following work of Zassenhaus.

The derangement subgroup

A few years ago, inspired by work of H. Zantema in number theory, Rosemary Bailey, Michael Giudici, Gordon Royle and I looked at the more general question. Let G be a finite transitive permutation group, and let $D(G)$ be the subgroup generated by the derangements in G . Which groups can arise as $G/D(G)$?

(This fits well our paradigm of inverse group theory.)

We found that the vast majority of transitive groups have $D(G) = G$. Of course, if G is a Frobenius group, then $D(G)$ is the Frobenius kernel, and the quotient is the Frobenius complement. The structure of Frobenius complements is known very precisely following work of Zassenhaus.

We wondered at first whether every derangement quotient is isomorphic to a Frobenius complement, but found a few examples where this was not the case.

The abstract to the rescue!

Unbekown to me (to my shame), Helmut Wielandt had proved a nice generalization of Frobenius' theorem, and thus defined what are called **Frobenius–Wielandt complements**, or **FW-complements**.

The abstract to the rescue!

Unbekown to me (to my shame), Helmut Wielandt had proved a nice generalization of Frobenius' theorem, and thus defined what are called **Frobenius–Wielandt complements**, or **FW-complements**.

Moreover, Carlo Scoppola had worked on these things, and come to a good understanding of which groups of prime power order could arise.

The abstract to the rescue!

Unbekown to me (to my shame), Helmut Wielandt had proved a nice generalization of Frobenius' theorem, and thus defined what are called **Frobenius–Wielandt complements**, or **FW-complements**.

Moreover, Carlo Scoppola had worked on these things, and come to a good understanding of which groups of prime power order could arise.

Last year, at the Ischia Group Theory Conference, I mentioned our work, and in later discussions with Carlo and (via him) Norberto Gavioli, we reached the conclusion that the two approaches were almost exactly equivalent (not quite, for rather technical reasons), but that Carlo's work gave us many more examples of derangement quotients which are p -groups.

The abstract to the rescue!

Unbekown to me (to my shame), Helmut Wielandt had proved a nice generalization of Frobenius' theorem, and thus defined what are called **Frobenius–Wielandt complements**, or **FW-complements**.

Moreover, Carlo Scoppola had worked on these things, and come to a good understanding of which groups of prime power order could arise.

Last year, at the Ischia Group Theory Conference, I mentioned our work, and in later discussions with Carlo and (via him) Norberto Gavioli, we reached the conclusion that the two approaches were almost exactly equivalent (not quite, for rather technical reasons), but that Carlo's work gave us many more examples of derangement quotients which are p -groups. The general problem of determining all derangement quotients is still open.



The ADE affair

This relates to a question raised by Vladimir Arnold as a “modern-day Hilbert problem” in an AMS symposium on the Hilbert problems in the 1970s.

The ADE affair

This relates to a question raised by Vladimir Arnold as a “modern-day Hilbert problem” in an AMS symposium on the Hilbert problems in the 1970s.

Lie algebras form a somewhat abstract subject. They are the structures which naturally live on the tangent space at the identity of a **Lie group**, a group which is also a manifold, such that the multiplication and inversion are continuous. These groups are widely used in theoretical physics, and they are best studied via their Lie algebras.

The ADE affair

This relates to a question raised by Vladimir Arnold as a “modern-day Hilbert problem” in an AMS symposium on the Hilbert problems in the 1970s.

Lie algebras form a somewhat abstract subject. They are the structures which naturally live on the tangent space at the identity of a **Lie group**, a group which is also a manifold, such that the multiplication and inversion are continuous. These groups are widely used in theoretical physics, and they are best studied via their Lie algebras.

The classification of finite-dimensional simple Lie algebras over the complex numbers was achieved by Cartan and Killing in the early 20th century. It turns out to reduce to a very concrete problem.

Root systems

A **Cartan subalgebra** of a Lie algebra is a nilpotent self-normalising subalgebra. For simple Lie algebras over \mathbb{C} , this subalgebra is abelian (the Lie bracket is trivial on it), but has a natural Euclidean inner product; it also contains a finite set of vectors called a *root system*, the roots being eigenvectors of the adjoint transformation induced by elements outside.

Root systems

A **Cartan subalgebra** of a Lie algebra is a nilpotent self-normalising subalgebra. For simple Lie algebras over \mathbb{C} , this subalgebra is abelian (the Lie bracket is trivial on it), but has a natural Euclidean inner product; it also contains a finite set of vectors called a *root system*, the roots being eigenvectors of the adjoint transformation induced by elements outside. The root system S has the properties

Root systems

A **Cartan subalgebra** of a Lie algebra is a nilpotent self-normalising subalgebra. For simple Lie algebras over \mathbb{C} , this subalgebra is abelian (the Lie bracket is trivial on it), but has a natural Euclidean inner product; it also contains a finite set of vectors called a *root system*, the roots being eigenvectors of the adjoint transformation induced by elements outside.

The root system S has the properties

- ▶ if $v \in S$, then $cv \in S$ if and only if $c = \pm 1$;

Root systems

A **Cartan subalgebra** of a Lie algebra is a nilpotent self-normalising subalgebra. For simple Lie algebras over \mathbb{C} , this subalgebra is abelian (the Lie bracket is trivial on it), but has a natural Euclidean inner product; it also contains a finite set of vectors called a *root system*, the roots being eigenvectors of the adjoint transformation induced by elements outside.

The root system S has the properties

- ▶ if $v \in S$, then $cv \in S$ if and only if $c = \pm 1$;
- ▶ for any $v, w \in S$, $2(v \cdot w) / (v \cdot v)$ is an integer (this is the **crystallographic condition**;

Root systems

A **Cartan subalgebra** of a Lie algebra is a nilpotent self-normalising subalgebra. For simple Lie algebras over \mathbb{C} , this subalgebra is abelian (the Lie bracket is trivial on it), but has a natural Euclidean inner product; it also contains a finite set of vectors called a *root system*, the roots being eigenvectors of the adjoint transformation induced by elements outside.

The root system S has the properties

- ▶ if $v \in S$, then $cv \in S$ if and only if $c = \pm 1$;
- ▶ for any $v, w \in S$, $2(v \cdot w) / (v \cdot v)$ is an integer (this is the **crystallographic condition**;
- ▶ the reflection in the hyperplane perpendicular to any one of its vectors maps S to itself.

Classification of root systems

The method I describe is not the original one but I think it is conceptually simpler.

Classification of root systems

The method I describe is not the original one but I think it is conceptually simpler.

It is clear from the definition that the roots of fixed length in a root system form a root system in their own right. So as a first step we assume all roots have the same length, which we take to be $\sqrt{2}$. Then the inner product of two independent roots is in $\{0, \pm 1\}$, so they make angles 90° , 60° or 120° .

Classification of root systems

The method I describe is not the original one but I think it is conceptually simpler.

It is clear from the definition that the roots of fixed length in a root system form a root system in their own right. So as a first step we assume all roots have the same length, which we take to be $\sqrt{2}$. Then the inner product of two independent roots is in $\{0, \pm 1\}$, so they make angles 90° , 60° or 120° .

Moreover, we can choose a basis such that all inner products are non-positive.

Classification of root systems

The method I describe is not the original one but I think it is conceptually simpler.

It is clear from the definition that the roots of fixed length in a root system form a root system in their own right. So as a first step we assume all roots have the same length, which we take to be $\sqrt{2}$. Then the inner product of two independent roots is in $\{0, \pm 1\}$, so they make angles 90° , 60° or 120° .

Moreover, we can choose a basis such that all inner products are non-positive.

The **Gram matrix** of inner products of this set thus has the form $2I - A$, where A is a symmetric 0, 1-matrix, hence the adjacency matrix of a graph whose greatest eigenvalue is less than 2.

ADE

Now we have an extremely concrete problem: find all the finite graphs (which we may assume connected) whose greatest eigenvalue is less than 2.

ADE


Now we have an extremely concrete problem: find all the finite graphs (which we may assume connected) whose greatest eigenvalue is less than 2.

This problem was only solved by J. H. Smith in 1969. The graphs are the **ADE diagrams**:

ADE

Now we have an extremely concrete problem: find all the finite graphs (which we may assume connected) whose greatest eigenvalue is less than 2.


This problem was only solved by J. H. Smith in 1969. The graphs are the **ADE diagrams**:



ADE

Now we have an extremely concrete problem: find all the finite graphs (which we may assume connected) whose greatest eigenvalue is less than 2.

This problem was only solved by J. H. Smith in 1969. The graphs are the **ADE diagrams**:



The other root systems (B_n , C_n , G_2 and F_4) can now be found by solving an easy puzzle involving putting together scaled copies of direct sums of these.

More

The ADE diagrams are ubiquitous in mathematics, involved in the **McKay correspondence** (connecting them with the rotation groups in Euclidean 3-space, viz. cyclic, dihedral, tetrahedral, octahedral and icosahedral groups),

More

The ADE diagrams are ubiquitous in mathematics, involved in the **McKay correspondence** (connecting them with the rotation groups in Euclidean 3-space, viz. cyclic, dihedral, tetrahedral, octahedral and icosahedral groups), **cluster algebras**,

More

The ADE diagrams are ubiquitous in mathematics, involved in the **McKay correspondence** (connecting them with the rotation groups in Euclidean 3-space, viz. cyclic, dihedral, tetrahedral, octahedral and icosahedral groups), **cluster algebras**, **singularity theory**,

More

The ADE diagrams are ubiquitous in mathematics, involved in the **McKay correspondence** (connecting them with the rotation groups in Euclidean 3-space, viz. cyclic, dihedral, tetrahedral, octahedral and icosahedral groups), **cluster algebras**, **singularity theory**, **algebras of finite representation type**,

More

The ADE diagrams are ubiquitous in mathematics, involved in the **McKay correspondence** (connecting them with the rotation groups in Euclidean 3-space, viz. cyclic, dihedral, tetrahedral, octahedral and icosahedral groups), **cluster algebras**, **singularity theory**, **algebras of finite representation type**, **locally flat spacetimes**, and so on.

More

The ADE diagrams are ubiquitous in mathematics, involved in the **McKay correspondence** (connecting them with the rotation groups in Euclidean 3-space, viz. cyclic, dihedral, tetrahedral, octahedral and icosahedral groups), **cluster algebras**, **singularity theory**, **algebras of finite representation type**, **locally flat spacetimes**, and so on. Arnold's question was to explain the correspondences between these many occurrences.

More

The ADE diagrams are ubiquitous in mathematics, involved in the **McKay correspondence** (connecting them with the rotation groups in Euclidean 3-space, viz. cyclic, dihedral, tetrahedral, octahedral and icosahedral groups), **cluster algebras**, **singularity theory**, **algebras of finite representation type**, **locally flat spacetimes**, and so on. Arnold's question was to explain the correspondences between these many occurrences.

My own first involvement with them was in the paper with Jean-Marie Goethals, Jaap Seidel, and Ernie Shult, where we settled a conjecture of Alan Hoffman by finding the graphs with least eigenvalue -2 . This is not the dual of graphs with greatest eigenvalue 2 (there are many such graphs), but the ADE structures are used in the proof (we also gave a new proof of the classification).

The book

At the time of his death a few years ago, John McKay and his research assistant Yang He were writing a book about all things ADE. Yang took over the project and recruited Pierre Dechant and me to join.

The book

At the time of his death a few years ago, John McKay and his research assistant Yang He were writing a book about all things ADE. Yang took over the project and recruited Pierre Dechant and me to join.

The book, *ADE: Patterns in Mathematics* will be published by Cambridge University Press in the London Mathematical Society Lecture Note Series later this month. (It is now available for pre-order.)

The book

At the time of his death a few years ago, John McKay and his research assistant Yang He were writing a book about all things ADE. Yang took over the project and recruited Pierre Dechant and me to join.

The book, *ADE: Patterns in Mathematics* will be published by Cambridge University Press in the London Mathematical Society Lecture Note Series later this month. (It is now available for pre-order.)

It is the first book in the series not to have a plain blue cover: the publisher and the LMS agreed to put the ADE diagrams on the cover.



Conclusion

The purpose of life is to prove and to conjecture.

Paul Erdős

Conclusion

The purpose of life is to prove and to conjecture.

Paul Erdős

I am now retired. This does not mean that I am giving up mathematics (of course!) But it is time to pass on the torch. So please continue: solve my problems; prove (or disprove) my conjectures; and, most important, have fun!



Conclusion

The purpose of life is to prove and to conjecture.

Paul Erdős

I am now retired. This does not mean that I am giving up mathematics (of course!) But it is time to pass on the torch. So please continue: solve my problems; prove (or disprove) my conjectures; and, most important, have fun!



for your attention!